

Sum rule for a difference of proton and neutron total photoproduction cross-sections

E. Bartoš,¹ S. Dubníčka,² and E. A. Kuraev¹

¹*Joint Institute for Nuclear Research, 141980 Dubna, Russia*

²*Inst. of Physics, Slovak Acad. of Sci., Dúbravská cesta 9, 845 11 Bratislava, Slovakia*

Starting from very high energy inelastic electron-nucleon scattering with a production of a hadronic state X to be moved closely to the direction of the initial nucleon, then utilizing analytic properties of parts of forward virtual Compton scattering amplitudes on proton and neutron, one obtains the relation between nucleon form factors and a difference of proton and neutron differential electroproduction cross-sections. In particular, for the case of small transferred momenta, one finally derives sum rule, relating Dirac proton mean square radius and anomalous magnetic moments of proton and neutron to the integral over a difference of the total proton and neutron photoproduction cross-sections.

PACS numbers: 11.55.Hx, 13.60.Hb, 25.20.Lj

Keywords: sum rule, photoproduction, cross-section

At the end of sixties of the last century Kurt Gottfried, by a consideration of the very high-energy electron-proton scattering and the nonrelativistic quark model of hadrons, has found [1]¹ a sum rule relating the proton mean square charge radius $\langle r_{Ep}^2 \rangle$ and the proton magnetic moment $\mu_p = 1 + \kappa_p$ to the integral over the total proton photoproduction cross-section $\sigma_{tot}^p(\nu)$ in the form

$$\int_0^\infty \frac{d\nu}{\nu} \sigma_{tot}^p(\nu) = \frac{\pi^2 \alpha}{m_p^2} \left[\frac{4}{3} m_p^2 \langle r_{Ep}^2 \rangle + 1 - \mu_p^2 \right], \quad (1)$$

where ν is the energy loss in the laboratory frame, α is the fine structure constant and m_p is the proton mass.

Nowadays it is well known, that the Gottfried sum rule cannot be satisfied since the corresponding integral diverges due to the known rise of the total proton photoproduction cross-section at high energies.

In this paper by means of a distinct way from the Gottfried approach and considering the nucleon isodoublet (proton and neutron) simultaneously a new sum rule is derived, which relates Dirac proton mean square radius and anomalous magnetic moments of proton and neutron to the integral over a difference of the total proton and neutron photoproduction cross-sections. Thus the rise of both photoproduction cross-sections at high energies is mutually cancelled and the corresponding integral converges.

In a derivation of the new sum rule we start also with a consideration of the very high energy electron-nucleon scattering

$$e^-(p_1) + N(p) \rightarrow e^-(p'_1) + X \quad (2)$$

with a production of a hadronic state X moving closely to the direction of initial nucleon. The corresponding one

photon exchange approximation matrix element takes the form

$$M = i \frac{\sqrt{4\pi\alpha}}{q^2} \bar{u}(p'_1) \gamma_\mu u(p_1) \langle X | J_\nu^{QED} | N^{(r)} \rangle g^{\mu\nu} \quad (3)$$

where (r) means a spin-state of the nucleon.

If the Gribov representation [2] of the metric tensor in the photon propagator of (3) is applied

$$g_{\mu\nu} = g_{\mu\nu}^\perp + \frac{2}{s} (\tilde{p}_\mu \tilde{p}_{1\nu} + \tilde{p}_\nu \tilde{p}_{1\mu}) \approx \frac{2}{s} \tilde{p}_\mu \tilde{p}_{1\nu}, \quad s = (p_1 + p)^2 \quad (4)$$

where

$$\tilde{p}_1 = p_1 - m_e^2 p / (2p_1 p), \quad \tilde{p} = p - m_N^2 p_1 / (2p_1 p) \quad (5)$$

are almost light-like vectors to be used according to Sudakov [3] for an expansion of the photon transferred four-momentum q

$$q = \beta_q \tilde{p}_1 + \alpha_q \tilde{p} + q_\perp, \quad q_\perp = (0, 0, \mathbf{q}), \quad (6)$$

$$\tilde{p} q_\perp = \tilde{p}_1 q_\perp = 0, \quad q_\perp^2 = -\mathbf{q}^2,$$

for the corresponding cross-section one obtains

$$d\sigma = \frac{4\pi\alpha}{s(q^2)^2} p_1^\mu p_1^\nu \times \sum_{X \neq N} \sum_{r=-1/2}^{1/2} \langle N^{(r)} | J_\mu^{QED} | X \rangle^* \langle X | J_\nu^{QED} | N^{(r)} \rangle d\Gamma, \quad (7)$$

where summations through the created hadronic states X and the spin states of the initial nucleon are carried out and $d\Gamma$ denotes the final state phase-space volume. Further, approximating square momentum of virtual photon

$$q^2 \approx -[\mathbf{q}^2 + (m_e s_1/s)^2] \quad (8)$$

¹ We would like to thank S. B. Gerasimov for bringing this paper to our attention.

with

$$s_1 = 2(qp) = s\beta_q, \quad (9)$$

i. e., it is related to the invariant square mass of the final hadronic state by the relation

$$m_X^2 = s_1 + q^2 + m_N^2, \quad (10)$$

and transforming the phase space volume of the final electron into the form

$$\frac{1}{(2\pi)^3} \frac{d^3 p'_1}{2\epsilon'_1} = \frac{1}{(2\pi)^3} d^4 q \delta[(p_1 - q)^2] = \frac{1}{(2\pi)^3} \frac{ds_1}{2s} d^2 q_\perp, \quad (11)$$

one gets the final state phase-space volume in the form

$$d\Gamma = \frac{ds_1}{2s(2\pi)^3} d^2 q_\perp d\Gamma_X; \quad (12)$$

$$d\Gamma_X = (2\pi)^4 \delta^4(q + p - \sum_j p_j) \prod_j \frac{d^3 p_j}{2\epsilon_j (2\pi)^3}.$$

Besides, using the current conservation condition

$$q^\mu \langle X | J_\mu^{QED} | N^{(r)} \rangle \approx (\beta_q p_1 + q_\perp)^\mu \langle X | J_\mu^{QED} | N^{(r)} \rangle = 0, \quad (13)$$

($\alpha_q \tilde{p}$ gives a negligible contribution) one can write

$$\begin{aligned} & \int p_1^\mu p_1^\nu \sum_{X \neq N} \sum_{r=-1/2}^{1/2} \langle N^{(r)} | J_\mu^{QED} | X \rangle^* \times \\ & \langle X | J_\nu^{QED} | N^{(r)} \rangle d\Gamma_X = p_1^\mu p_1^\nu \Delta_{\mu\nu}^{(N)} = \\ & \frac{s^2}{s_1^2} q^2 e^i e^j \Delta_{ij}^{(N)} = 2i \frac{s^2}{s_1^2} q^2 \text{Im} \tilde{A}^{(N)}(s_1, \mathbf{q}), \quad e_i = \frac{\mathbf{q}_i}{|\mathbf{q}|}, \end{aligned} \quad (14)$$

where just Cutkosky rules for s -channel discontinuity $\Delta \tilde{A}^{(N)} = 2i \text{Im} \tilde{A}^{(N)}$ of the corresponding Feynman amplitude was applied.

Here we would like to note that the amplitude $\tilde{A}(s_1, \mathbf{q})$ by a construction is only a part of the total forward virtual Compton scattering amplitude $A(s_1, \mathbf{q})$, which doesn't contain (unlike the amplitude $A(s_1, \mathbf{q})$) any crossing Feynman diagram contributions. As a result there is no u -channel pole in $\tilde{A}(s_1, \mathbf{q})$, which is a crucial point in a derivation of the new sum rule by using the analytic properties of the latter amplitude.

Since the imaginary part of the crossing Feynman diagrams is starting to be different from zero only above

$$s_1^{(3N)} = 8m_N^2 + \mathbf{q}^2, \quad (15)$$

one can write down an equality relation

$$\text{Im} \tilde{A}(s_1, \mathbf{q}) = \text{Im} A(s_1, \mathbf{q}) = 4s_1 \sigma_{tot}^{\gamma^* p \rightarrow X}(s_1, \mathbf{q}), \quad (16)$$

for $2m_N m_\pi + m_\pi^2 + \mathbf{q}^2 \leq s_1 \leq 8m_N^2 + \mathbf{q}^2$. Fortunately above the threshold (15) total proton and neutron photoproduction cross-sections are almost equal and at the new sum rule one can integrate over them up to infinity.

Now, substituting relation (14) into cross-section expression (7) and integrating over the phase space volume of the final hadronic state X , as well as over the invariant mass squared m_X^2 , i. e., over the variable s_1 (see (10)), to be interested only for $\mathbf{q}$ distribution, for a difference of corresponding differential proton and neutron electroproduction cross-sections one finds (see Appendix D in [4])

$$\begin{aligned} & \left(\frac{d\sigma^{e^- p \rightarrow e^- X}(s, \mathbf{q})}{d^2 \mathbf{q}} - \frac{d\sigma^{e^- n \rightarrow e^- X}(s, \mathbf{q})}{d^2 \mathbf{q}} \right) = \\ & \frac{\alpha q^2}{4\pi^2} \int_{2m_N m_\pi + m_\pi^2 + \mathbf{q}^2}^{\infty} \frac{ds_1}{s_1^2 [q^2 + (m_e s_1/s)^2]^2} \times \\ & \left(\text{Im} \tilde{A}^{(p)}(s_1, \mathbf{q}) - \text{Im} \tilde{A}^{(n)}(s_1, \mathbf{q}) \right). \end{aligned} \quad (17)$$

If one neglects the second term in square brackets of the denominator of the integral of (17) (due to the small value of m_e and high s in comparison with s_1) and takes into account (16) for $\mathbf{q}^2 \rightarrow 0$ together with the relation $d^2 \mathbf{q} = \pi d\mathbf{q}^2$ one comes to the expression

$$\begin{aligned} & q^2 \left(\frac{d\sigma^{e^- p \rightarrow e^- X}}{d\mathbf{q}^2} - \frac{d\sigma^{e^- n \rightarrow e^- X}}{d\mathbf{q}^2} \right) \Big|_{\mathbf{q}^2 \rightarrow 0} = \\ & \frac{\alpha}{\pi} \int_{2m_N m_\pi + m_\pi^2}^{\infty} \frac{ds_1}{s_1} \left(\sigma_{tot}^{\gamma p \rightarrow X}(s_1) - \sigma_{tot}^{\gamma n \rightarrow X}(s_1) \right), \end{aligned} \quad (18)$$

similar to the difference of the total cross-sections of the electroproduction processes on proton and neutron in the Weizsäcker-Williams approximation [5, 6]

$$\begin{aligned} & \left(\sigma_{tot}^{e^- p \rightarrow e^- X}(s) - \sigma_{tot}^{e^- n \rightarrow e^- X}(s) \right) = \\ & 2 \frac{\alpha}{\pi} \ln \left(\frac{s}{m_e m_\pi} \right) \times \\ & \int_{2m_N m_\pi + m_\pi^2}^{\infty} \frac{ds_1}{s_1} \left(\sigma_{tot}^{\gamma p \rightarrow X}(s_1) - \sigma_{tot}^{\gamma n \rightarrow X}(s_1) \right). \end{aligned} \quad (19)$$

The analytic properties of the amplitude $\tilde{A}(s_1, \mathbf{q})$ in s_1 -plane are consisting of the one-nucleon intermediate state pole at $s_1 = \mathbf{q}^2$, the right-hand cut starting at the pion-nucleon threshold $s_1 = 2m_N m_\pi + m_\pi^2 + \mathbf{q}^2$ and the left-hand cut starting from $s_1 = -\mathbf{q}^2 - 8m_N^2$.

If one defines the path integral I in s_1 -plane as presented in Fig. 1a

$$I = \int_C ds_1 \frac{p_1^\mu p_1^\nu}{s^2} \left(\tilde{A}_{\mu\nu}^{(p)}(s_1, \mathbf{q}) - \tilde{A}_{\mu\nu}^{(n)}(s_1, \mathbf{q}) \right) \quad (20)$$




Fig. 1: Sum rule interpretation in s_1 plane. On subfigure (a) is drawn the contour C [see (20)], on the subfigure (b) is the contour C closed to upper and lower half plane.

then once the contour C is closed to upper half-plane, another one to lower half-plane (Fig. 1b) and considering (14), the following sum rule

$$\pi \left(\text{Res}^{(n)} - \text{Res}^{(p)} \right) = \mathbf{q}^2 \int_{r.h.}^{\infty} \frac{ds_1}{s_1^2} \left(\text{Im} \tilde{A}^{(p)}(s_1, \mathbf{q}) - \text{Im} \tilde{A}^{(n)}(s_1, \mathbf{q}) \right) \quad (21)$$

appears with (an averaging through the initial nucleon and photon spins is performed)

$$\text{Res}^{(N)} = 2\pi\alpha \left(F_{1N}^2 + \frac{\mathbf{q}^2}{4m_N^2} F_{2N}^2 \right) \quad (22)$$

to be the one-nucleon intermediate state pole contribution and the left-hand (*l.h.*) cut contributions from the difference $\left(\text{Im} \tilde{A}^{(p)} - \text{Im} \tilde{A}^{(n)} \right)$ are mutually annulated.

Substituting (22) into (21) and taking into account (17) one comes to the relation

$$F_{1n}^2(-\mathbf{q}^2) + \frac{\mathbf{q}^2}{4m_n^2} F_{2n}^2(-\mathbf{q}^2) - F_{1p}^2(-\mathbf{q}^2) - \frac{\mathbf{q}^2}{4m_p^2} F_{2p}^2(-\mathbf{q}^2) = 2 \frac{(\mathbf{q}^2)^2}{\pi\alpha^2} \left(\frac{d\sigma^{e^-p \rightarrow e^-X}}{d\mathbf{q}^2} - \frac{d\sigma^{e^-n \rightarrow e^-X}}{d\mathbf{q}^2} \right). \quad (23)$$

For $\mathbf{q}^2 = 0$ the right-hand side is equal to zero, but the left-hand side is -1 . This is caused by the following reasons. On the right-hand side we take into account only strong interactions effects and on the left-hand side the -1 is given by nonzero proton charge, i. e., pure electromagnetic effect. In order to separate the pure strong interactions from electromagnetic ones on the left-hand side of the sum rule one has to regularize the latter by

adding $+1$ in order to achieve the zero also there. As a result the sum rule takes the final form

$$1 + F_{1n}^2(-\mathbf{q}^2) + \frac{\mathbf{q}^2}{4m_n^2} F_{2n}^2(-\mathbf{q}^2) - F_{1p}^2(-\mathbf{q}^2) - \frac{\mathbf{q}^2}{4m_p^2} F_{2p}^2(-\mathbf{q}^2) = 2 \frac{(\mathbf{q}^2)^2}{\pi\alpha^2} \left(\frac{d\sigma^{e^-p \rightarrow e^-X}}{d\mathbf{q}^2} - \frac{d\sigma^{e^-n \rightarrow e^-X}}{d\mathbf{q}^2} \right), \quad (24)$$

giving into a relation nucleon electromagnetic form factors with a difference of the differential cross-sections of deep inelastic electron-proton scattering. There is a challenge to specialized experimental groups (e. g., in DESY) to verify the sum rule (24).

Now, taking a derivative of both sides in (24) according to $\mathbf{q}^2$ for $\mathbf{q}^2 \rightarrow 0$ and employing the relation (18) one comes to the new sum rule relating Dirac proton mean square radius and anomalous magnetic moments of proton and neutron

$$\langle r_{1p}^2 \rangle = 6 \frac{d}{d\mathbf{q}^2} F_{1p}(q^2) \Big|_{q^2=0}, \quad \kappa_N = F_{2N}(q^2) \Big|_{q^2=0} \quad (25)$$

to the integral over a difference of the total proton and neutron photoproduction cross-sections (we used for laboratory frame $s_1 = 2M_N\omega$)

$$\frac{1}{3} \langle r_{1p}^2 \rangle - \frac{\kappa_p^2}{4m_p^2} + \frac{\kappa_n^2}{4m_n^2} = \frac{2}{\pi^2\alpha} \int_{\omega_N}^{\infty} \frac{d\omega}{\omega} \left[\sigma_{tot}^{\gamma p \rightarrow X}(\omega) - \sigma_{tot}^{\gamma n \rightarrow X}(\omega) \right] \quad (26)$$

with $\omega_N = m_\pi + m_\pi^2/2M_N$, in which just a mutual cancellation of the rise of these total proton and neutron photoproduction cross-sections for $\omega \rightarrow \infty$, created by the Pomeron exchanges, is achieved. It is straightforward to see that this final sum rule is not influenced at

all by a renormalization of the left-hand side in (24) by +1.

Using the Dirac proton mean square radius from [7] and proton and neutron anomalous magnetic moments from [8], the evaluation of the left-hand side in (26) gives (1.93 ± 0.18) mbarn.

On the other hand the data on photoproduction cross section on the neutron are not so known as for the proton case up to now. Nevertheless, taking compilation of both of them from [9], and assuming that both total cross-sections are starting at the pion mass and are equal above the last neutron experimental point at $\omega = 17.84$ GeV one gets on the right-hand side the value (1.92 ± 0.32) mbarn. So, the sum rule (26) can be considered to be satisfied.

Acknowledgments

We are grateful to Lev Lipatov for valuable discussions and participants of the seminar at BLTP JINR, Dubna for comments. One of the authors (E.A.K) is grateful to ECT (Trento) Centrum. E.A.K and E.B would like to thank to Institute of Physics SAS in Bratislava for a warm hospitality.

The work was in part supported by Slovak Grant Agency for Sciences, Gr. No. 2/1111/23, E.A.K. is grateful to INTAS grant 00366.

-
- [1] K. Gottfried, Phys. Rev. Lett. **18** (1967) 1174.
 - [2] V. N. Gribov: Lectures on the theory of complex angular momenta, Phys.-Tech. Institute, Kharkov (1970).
 - [3] V. V. Sudakov, Sov. Phys. JETP **3** (1956) 65.
 - [4] V. N. Baier, V. S. Fadin, V. A. Khoze and E. A. Kuraev, Phys. Rep. **78** (1981) 293.
 - [5] G. Weizsäcker, Z. Phys. **88** (1934) 612.
 - [6] E. Williams, Phys. Rev. **45** (1939) 729.
 - [7] S. Dubnicka, A. Z. Dubnickova and P. Weisenpacher, J. Phys. **G 29** (2003) 405.
 - [8] K. Hagiwara et al, Phys. Rev. **D 66** (2002) 010001.
 - [9] A. Baldini et al, in Group I Elementary Particles, Nuclei and Atoms, edited by H. Schopper, Landolt-Börnstein, New Series, Vol. 12 (Springer-Verlag, Berlin, 1988).